

Bose-Einstein condensates in optical lattices

Creating number squeezed states of atoms

Matthew Davis

University of Queensland

Overview

- What is a BEC?
- What is an optical lattice?
- What happens to a BEC in an optical lattice?
- Experiments
 - Mott insulator
 - Collapse and revival of matter waves
- Other interesting bits
- Summary

Bose-Einstein condensation

- Definition: Macroscopic occupation of a single quantum level (usually ground state).
- Due to *saturation* of excited states of system


$$\text{Condition : } n\Lambda_{dB}^3 \geq 2.612$$

- Created for the first time in 1995 in atomic Rb
- More than 30 groups have now observed BEC
- Nobel prize for Cornell, Ketterle and Wieman in 2001.

BEC features

- Made by first laser cooling, then evaporative cooling a sample of trapped atoms.
- Occurs at ultracold temperatures ~ 100 nK.
- Typically consist of 10^4 – 10^7 atoms of eg Rb, Na.
- Best thought of as a *matter wave*, analogous to laser light.
- All atoms are phase coherent i.e. widely spaced atoms can form an interference pattern.
- *Quantum mechanics on a macroscopic scale*

Phase transition



Optical dipole force

- Stark effect: Shift of atomic energy levels in static electric field.
- AC Stark effect: Shift of atomic energy levels due to time-varying electric field of laser.
- In the limit of large detuning from resonance (avoid spontaneous decay)

$$\Delta E(x) \propto \frac{I(x)}{(\omega - \omega_0)^2}$$


- Remember $F(x) = -dE(x)/dx$, so atoms in a laser beam experience force proportional to gradient of intensity of laser beam

Optical lattices

- Definition : Standing waves of light
- Atoms see series of potential wells (hills).
- If cold enough, atoms can be trapped in single well.
- Blue-detuned : atoms trapped at nodes
- Red-detuned : atoms trapped at anti-nodes
- *Depth of potential easily controlled via laser intensity*

Loading BEC into lattice

- Increase intensity adiabatically
- $V(x) = E_R(t) \cos^2(kx/2), \quad k = 2\pi/\lambda$



The physics

Initially ...

- Atoms can tunnel between wells
- Each well is approximately a coherent state.
- Variance in atom number is equal to mean.
- System is superfluid.

As laser intensity increases ...

- Wells become isolated (longer tunnelling time).
- Number fluctuations suppressed.
- Heisenberg says — *phase fluctuations increase*.

Quantum phase transitions



- Classical phase transitions driven by thermal fluctuations.
- Quantum phase transitions driven by quantum fluctuations.
- Occur at $T=0$ by changing parameters of Hamiltonian.
- Mott insulator : Energy minimised by number states in each well.
- How? — reduce tunnelling rate by increasing lattice depth.

Experiments

1D lattice expt by Orzel et al, Science **291**, 2386 (2001).


3D lattice ext by Greiner et al, Nature **415**, 39 (2002).

Turn on lattice adiabatically, then switch off suddenly




Superfluid to Mott insulator ...

Images after 18 ms of expansion



... and back to superfluid




Adiabatic ramp to
 $22 E_R$ (80 ms)

Hold (20 ms)

Release to $9 E_R$
(t ms)

... and back to superfluid II



Theory: Bose-Hubbard model

$$\hat{H} = -J \sum_{\langle i,j \rangle} \hat{b}_i^\dagger \hat{b}_j + \sum_i \epsilon_i \hat{n}_i + \frac{U}{2} \sum_i \hat{n}_i (\hat{n}_i - 1)$$

J : tunneling matrix element. Decreases with E_R .

U : interaction matrix element. Increases with E_R .

Mott insulator transition: $\frac{U}{nJ} > 5.8 \times z$

n : number of atoms per site

z : number of nearest neighbours

More theory

- Recent results show that disappearance of interference is not a signature of MI phase (Roth and Burnett, condmat/0209066).
- However excitation spectrum was also studied.
- Energy gap appears above lattice depth of $13 E_{\text{recoil}}$.
- Thus an insulating state was achieved.

Collapse and revival

Coherent state: superposition of number states

$$|\alpha\rangle = e^{-|\alpha|^2/2} \sum_n \frac{\alpha^n}{\sqrt{n!}} |n\rangle$$

Ground state Hamiltonian of single well


$$\hat{H} = \frac{U}{2} \sum_i \hat{n}_i (\hat{n}_i - 1)$$

Evolution:


$$|\alpha\rangle(t) = e^{-|\alpha|^2/2} \sum_n \frac{\alpha^n}{\sqrt{n!}} e^{-i\frac{1}{2}Un(n-1)t/\hbar} |n\rangle$$

Revival

- All phases match up at times $t_c = U/h$



- Can be realised by sudden switch to Mott insulator state!




Increasing times in the MI state

Time scales

Revival time is *independent* of atom number variance.

Collapse time is *dependent* on initial variance of atom number



Other interesting bits

- Proposal (and experiments) on controlled cold collisions
Two different internal states, each see different lattice.
Move lattices together — interactions dependent on state.
Quantum computing???
- Proposal to form Mott insulator with $n = 2$
Then photoassociate and melt $\Rightarrow$ molecular BEC.
- Why is this interesting?
Wednesday physics seminar tomorrow, 1pm Conference room 2.32D, Physics Annexe (Building 6).
PAIR CORRELATED ATOM LASER BEAMS
Dr Karen Kheruntsyan.

Summary

- Introduction to BEC
- Introduction to optical lattices
- Quantum phase transitions
- Superfluid to Mott insulator
- Bose-Hubbard model.
- Collapse and revival of matter waves
- Other proposals for BECs in lattices.

No Microsoft products were used in the making of these slides

Thanks to linux, latex, prosper, and ps2pdf.